ZOOLOGY

time because the					
1) patients do not feel much pa	in				
	2) filarial worms enter into blood from lymph during night time				
3) microfilariae exhibit noctu					
	ood circulation during night time				
	a sense of euphoria and increased energy levels as it				
1)acts as amphetamine	2)stimulates adrenal glands to produce steroid				
3) causes the inhibition of chol	inergic nerves 4)interferes with the transport of dopamine				
38. Which of the following is no	t a symptom of chronic malaria?				
1)Splenomegaly	2)Hypertrophy of RBC				
3)Cirrhosis of liver	4)Anaemia				
39. In cockroach, the structures	that help in locomotion on smooth surface are				
1)claws and arolium	2) pulvillus and bristles of podomeres				
3)plantulae on tarsus	4)plantulae, claws				
40. Incorrect statement about th	e mushroom shaped gland of <i>Periplaneta americana</i> from				
the following					
1)Helps in storage and excretion	on of uric acid				
2)Secretes the innermost layer	of spermatophore				
3)Stores the sperms in the form	3)Stores the sperms in the form of spermatophores				
4)Nourishes the sperms					
41. In a biological community (Sauses principle explains				
1) competition among differen	1) competition among different species under unlimited resources				
2)competitive exclusion of an	inferior species due to the limited natural resources				
3) coexistence of closely relate	ed species due to resource partitioning				
4) intra specific competition du	ue to limited natural resources				
42. In India, grassland ecosyster	ns are found mainly in				
1)Ladakh region	2)Himalayan region				
3)Deccan plateau	4)Western ghats				
ROUGH					

143.	The influence of light on non-directional movement of animals is known as				
	1)phototaxis	2)photoperiodism	3)photokinesis	4)phototropism	
144.	According to the st	udy of molecular sys	stematics, the kingdo	om replaced by the two	
	kingdoms bacteria a	and archebacteria is			
	1)Fungi	2)Protista	3)Monera	4)Metaphyta	
145.	Identify the correct	match from the folle	owing		
	1)Earth summit-Joha	nnesburg, South Africa	l		
	2)Biosphere reserves	-Norman Myers			
	3)Jim Corbett Nation	al park-first Indian nat	ional park in Gujarat		
	4)Cryopreservation-e	ex-situ conservation			
146.	In humans mucous	connective tissue is p	resent mainly in the		
	1)mucosa lining of gu	ıt	2)lining of alveoli of	lungs	
	3)umbilical chord of	newly born infants			
	4) walls of trachea, br	onchi and bronchioles			
147.	Cardiac muscles are	e highly resistant to f	fatigue mainly due to	the presence of	
	1)intercalated discs	2)branched muscle fi	bres and communication	on junctions	
	3)specialised autorhy	thmic muscle fibres			
	4)numerous sarcosor	nes and myoglobin mo	olecules		
148.	148. Identify the correct combination from the following				
	Scientific name Respiratory organs Excretory organs				
	1)Astacus	Dermal branchiae	Green glands		
	2)Aplysia	Ctenidia	Meta nephridia		
	3)Aphrodite	Parapodia	Protonephridia		
	4)Ascidian	Pharyngeal gill slits	Proboscis gland		
149.	9. Echinoderm with elongated body and without arms, spines is				
	1)Gorgonocephalus2)Thyone3)Clypeaster4)Antedon				
150.). Incorrect statement about "great white shark" from the following				
	1)Sharks exhibit physiological uraemia				
	2)They have placoid scales and heteroceracal caudal fin				
	3)Fertilization internal and viviparous in nature				
	4)Lobe finned fishes	with air bladder acting	as hydrostatic organ		
		ROUG	θH		
1					

151. Identify the correct match from the following				
101.	1)Gavialis-Vascula		2) <i>Corvus</i> -Urinar	v bladder present
	3) <i>Cervus</i> -Intra abc			cervical vertebrae
152	,		,	cervical vertebrae
132.	-	otelic animal from the	_	4)Comol
152	1)Land leech	2)Land snail	3)Desert fox	4)Camel
155.		from the following	2) C - 1	
	1)Paneth cells-lyso	•	2)Schawnn cells-	•
154	3)Dendritic cells-a		4)Mast cells-infla	ammation
154.		owing is an exotic car	-	
	1)Tunas	2)Common carp	3)Cat fish	4)Oil sardine
155.	Follicular hyperk			
	1)sex influenced tr		2)holandric chara	
	3)sexlinked domin		4)autosomal rece	ssive character
156.		st is developed mainly		
	1)confirm the preg	•	2) determin the se	
	,	osomal abnormalities		rect position of foetus
157.		owing disorder is not o		(2n+1) in human beings?
	1)Down syndrome		2)Turner's syndro	
	3)Kline felter sync		4)Patau syndrome	
158.	158. In a family, blood groups of children are entirely different from that of parents only if			
	the blood groups	of the parents are		
	1)A and B	2)AB and AB	3)AB and O	4)O and A or B
159.	Correct match fr	om the following		
	1)Mutation theory-	Bateson	2)Survival of the	fittest-Herbert Spencer
	3)Principles of geo	ology-Alfred Wallace	4)Biochemical re	capitulation -Ernst Haeckel
160. The first human like being frugivorous in habit and hunted with sharp stony objects				
	was named			
	1)Australopithecu	S	2)Homo neander	rthalensis
	3)Homo habilis		4)Homo erectus	
		ROU	GH	

1)Edward Jenner 2)Alexander Fleming 3)Louis Pasteur 4)Jonas Salk 163. Ampulla of vater is formed by the union of 1)common hepatic duct and pancreatic duct 2)common bile duct and pancreatic duct 3)cystic duct and common hepatic duct 4)infundibulum and isthmus of fallopian to 1.10.000000000000000000000000000000000	161.	Incorrect match fro	om the following			
 162. The technique of attenuating or weakening of a microbe was developed by Edward Jenner Alexander Fleming Jouis Pasteur Jonas Salk 163. Ampulla of vater is formed by the union of common hepatic duct and pancreatic duct common hepatic duct and pancreatic duct common hepatic duct and pancreatic duct a) infurnition of pancreatic duct a) inferior venacava a) inferior venacava b) infuri diventricular agenture 		1)MRI-aneurysm		2)EEG-epilepsy		
1)Edward Jenner 2)Alexander Fleming 3)Louis Pasteur 4)Jonas Salk 163. Ampulla of vater is formed by the union of 1)common hepatic duct and pancreatic duct 2)common bile duct and pancreatic duct 3)cystic duct and common hepatic duct 4)infundibulum and isthmus of fallopian to 1.10.000000000000000000000000000000000		3)PSA test-cervical c	ancer	4)PCR-HIV detection	on	
 163. Ampulla of vater is formed by the union of l)common hepatic duct and pancreatic duct c)common hepatic duct and pancreatic duct c)common hepatic duct and common hepatic duct d)infundibulum and isthmus of fallopian to 164. Adenoids are present in n)nasopharynx c)oropharynx 105. In human development, most of the organ systems are developed by the end of l)first month l)24 weeks l)first trimester weeks 106. The cells that provide nutrition to developing spermatozoa in testes are called l)Kupffer cells l)Leydig cells s)Sertoli cells d)Oxyntic cells 107. Rima glottidis is the narrow space between the l)true vocal folds and false vocal cords c)true vocal cords and arytenoids c)corniculate cartilages and cuneiform cartilages l)acterium Streptococcus pneumoniae l)inhalation of coal dust and sulphur particles release of vasodilator substances due to allergens d)deposition of iron particles in alveoli of lungs 106. The specialised myocytes present in inner ventricular walls of heart ben endocardium are called l)Chorde tendinae Papillary muscles Purkinje fibres Columnae carnea l)coronary sinus l)superior venacava l)inferior venacava 	162.	The technique of at	tenuating or weakeni	ng of a microbe wa	s developed by	
1)common hepatic duct and pancreatic duct 2)common bile duct and pancreatic duct 3)cystic duct and common hepatic duct 4)infundibulum and isthmus of fallopian to 10, 10, 10, 10, 10, 10, 10, 10, 10, 10,		1)Edward Jenner	2)Alexander Fleming	3)Louis Pasteur	4)Jonas Salk	
 3)cystic duct and common hepatic duct 4)infundibulum and isthmus of fallopian to 10,00000000000000000000000000000000000	163.	Ampulla of vater is	formed by the union	of		
164. Adenoids are present in 1)nasopharynx 2)oropharynx 3)below the tongue 4)laryngophrynx 165. In human development, most of the organ systems are developed by the end of 1)first month 2)24 weeks 3)below the tongue 4)laryngophrynx 166. The cells that provide nutrition to developing spermatozoa in testes are called 1)Kupffer cells 2)Leydig cells 3)bertoli cells 4)Oxyntic cells 107. Rima glottidis is the narrow space between the 1)true vocal folds and false vocal cords 2)true vocal cords and arytenoids 3)corniculate cartilages and cuneiform cartilages 4)arytenoids and thyroid cartilages 168. Pneumoconiosis is caused by the 1)bacterium Streptococcus pneumoniae 2)inhalation of coal dust and sulphur particles 3)release of vasodilator substances due to allergens 4)deposition of iron particles in alveoli of lungs 10 169. The specialised myocytes present in inner ventricular walls of heart ben endocardium are called 1)Chorde tendinae 1)Chorde tendinae 2)Papillary muscles 3)Purkinje fibres 4)Columnae carnea 170. In human heart, the non functional valve in the right atrium is present at the ope of 1)coronary sinus 2)superior venacava 3)inferior venacava		1)common hepatic du	ict and pancreatic duct	2)common bile duc	t and pancreatic duct	
1)nasopharynx 2)oropharynx 3)below the tongue 4)laryngophrynx 165. In human development, most of the organ systems are developed by the end of 1)first month 2)24 weeks 3)first trimester 4)8 weeks 166. The cells that provide nutrition to developing spermatozoa in testes are called 1)Kupffer cells 2)Leydig cells 1)Kupffer cells 2)Leydig cells 3)Sertoli cells 1)true vocal folds and false vocal cords 2)true vocal cords and arytenoids 3)corniculate cartilages and cuneiform cartilages 4)arytenoids and thyroid cartilages 108. Pneumoconiosis is caused by the 1)bacterium Streptococcus pneumoniae 2)inhalation of coal dust and sulphur particles 3)release of vasodilator substances due to allergens 4)deposition of iron particles in alveoli of lungs 169. The specialised myocytes present in inner ventricular walls of heart ben endocardium are called 1)Chorde tendinae 2)Papillary muscles 3)Purkinje fibres 4)Columnae carnea 170. In human heart, the non functional valve in the right atrium is present at the ope of 1)coronary sinus 2)superior venacava 3)inferior venacava 4)right atrioventricular aperture		3)cystic duct and con	nmon hepatic duct	4)infundibulum and	l isthmus of fallopian tube	
 3)below the tongue 4)laryngophrynx 165. In human development, most of the organ systems are developed by the end of 1)first month 2)24 weeks 3)first trimester 4)8 weeks 166. The cells that provide nutrition to developing spermatozoa in testes are called 1)Kupffer cells 2)Leydig cells 3)Sertoli cells 4)Oxyntic cells 1)Kupffer cells 2)Leydig cells 3)Sertoli cells 4)Oxyntic cells 4)Oxyntic cells 4)Corniculate cartilages and cuneiform cartilages 4)deposition of icon particles in alveoli of lungs 106. The specialised myocytes present in inner ventricular walls of heart ben endocardium are called 1)Chorde tendinae 2)Papillary muscles 3)Purkinje fibres 4)Columnae carnea 1)coronary sinus 2)superior venacava 3)inferior venacava 4)right atrioventricular aperture 	164.	Adenoids are presen	nt in			
 165. In human development, most of the organ systems are developed by the end of 1)first month 2)24 weeks 3)first trimester 4)8 weeks 166. The cells that provide nutrition to developing spermatozoa in testes are called 1)Kupffer cells 2)Leydig cells 3)Sertoli cells 4)Oxyntic cells 167. Rima glottidis is the narrow space between the 1)true vocal folds and false vocal cords 2)true vocal cords and arytenoids 3)corniculate cartilages and cuneiform cartilages 4)arytenoids and thyroid cartilages 168. Pneumoconiosis is caused by the 1)bacterium <i>Streptococcus pneumoniae</i> 2)inhalation of coal dust and sulphur particles 3)release of vasodilator substances due to allergens 4)deposition of iron particles in alveoli of lungs 169. The specialised myocytes present in inner ventricular walls of heart ben endocardium are called 1)Chorde tendinae 2)Papillary muscles 3)Purkinje fibres 4)Columnae carnea 170. In human heart, the non functional valve in the right atrium is present at the ope of 1)coronary sinus 2)superior venacava 3)inferior venacava 4)right atrioventricular aperture 		1)nasopharynx		2)oropharynx		
1)first month 2)24 weeks 3)first trimester 4)8 weeks 166. The cells that provide nutrition to developing spermatozoa in testes are called 1)Kupffer cells 2)Leydig cells 3)Sertoli cells 4)Oxyntic cells 167. Rima glottidis is the narrow space between the 1)true vocal folds and false vocal cords 2)true vocal cords and arytenoids 3)corniculate cartilages and cuneiform cartilages 4)arytenoids and thyroid cartilages 168. Pneumoconiosis is caused by the 1)bacterium Streptococccus pneumoniae 2)inhalation of coal dust and sulphur particles 3)release of vasodilator substances due to allergens 4)deposition of iron particles in alveoli of lungs 169. The specialised myocytes present in inner ventricular walls of heart ben endocardium are called 1)Chorde tendinae 2)Papillary muscles 3)Purkinje fibres 4)Columnae carnea 170. In human heart, the non functional valve in the right atrium is present at the ope of 1)coronary sinus 2)superior venacava 3)inferior venacava 4)right atrioventricular aperture		3) below the tongue		4)laryngophrynx		
 166. The cells that provide nutrition to developing spermatozoa in testes are called 1)Kupffer cells 2)Leydig cells 3)Sertoli cells 4)Oxyntic cells 167. Rima glottidis is the narrow space between the 1)true vocal folds and false vocal cords 2)true vocal cords and arytenoids 3)corniculate cartilages and cuneiform cartilages 4)arytenoids and thyroid cartilages 168. Pneumoconiosis is caused by the 1)bacterium <i>Streptococcus pneumoniae</i> 2)inhalation of coal dust and sulphur particles 3)release of vasodilator substances due to allergens 4)deposition of iron particles in alveoli of lungs 169. The specialised myocytes present in inner ventricular walls of heart ben endocardium are called 1)Chorde tendinae 2)Papillary muscles 3)Purkinje fibres 4)Columnae carnea 170. In human heart, the non functional valve in the right atrium is present at the ope of 1)coronary sinus 2)superior venacava 3)inferior venacava 	165.	In human developm	ent, most of the organ	n systems are devel	oped by the end of	
 167. Rima glottidis is the narrow space between the 1)true vocal folds and false vocal cords 2)true vocal cords and arytenoids 3)corniculate cartilages and cuneiform cartilages 4)arytenoids and thyroid cartilages 168. Pneumoconiosis is caused by the 1)bacterium <i>Streptococcus pneumoniae</i> 2)inhalation of coal dust and sulphur particles 3)release of vasodilator substances due to allergens 4)deposition of iron particles in alveoli of lungs 169. The specialised myocytes present in inner ventricular walls of heart ben endocardium are called 1)Chorde tendinae 2)Papillary muscles 3)Purkinje fibres 4)Columnae carnea 170. In human heart, the non functional valve in the right atrium is present at the ope of 1)coronary sinus 2)superior venacava 4)right atrioventricular aperture 		1)first month	2)24 weeks	3)first trimester	4)8 weeks	
 167. Rima glottidis is the narrow space between the 1)true vocal folds and false vocal cords 2)true vocal cords and arytenoids 3)corniculate cartilages and cuneiform cartilages 4)arytenoids and thyroid cartilages 168. Pneumoconiosis is caused by the 1)bacterium <i>Streptococcus pneumoniae</i> 2)inhalation of coal dust and sulphur particles 3)release of vasodilator substances due to allergens 4)deposition of iron particles in alveoli of lungs 169. The specialised myocytes present in inner ventricular walls of heart ben endocardium are called 1)Chorde tendinae 2)Papillary muscles 3)Purkinje fibres 4)Columnae carnea 170. In human heart, the non functional valve in the right atrium is present at the ope of 1)coronary sinus 2)superior venacava 4)right atrioventricular aperture 	166.	The cells that prov	ide nutrition to deve	loping spermatozoa	a in testes are called	
 3)corniculate cartilages and cuneiform cartilages 4)arytenoids and thyroid cartilages 168. Pneumoconiosis is caused by the 1)bacterium Streptococcus pneumoniae 2)inhalation of coal dust and sulphur particles 3)release of vasodilator substances due to allergens 4)deposition of iron particles in alveoli of lungs 169. The specialised myocytes present in inner ventricular walls of heart ben endocardium are called 1)Chorde tendinae 2)Papillary muscles 3)Purkinje fibres 4)Columnae carnea 170. In human heart, the non functional valve in the right atrium is present at the ope of 1)coronary sinus 2)superior venacava 4)right atrioventricular aperture 		1)Kupffer cells	2)Leydig cells	3)Sertoli cells	4)Oxyntic cells	
 3)corniculate cartilages and cuneiform cartilages 4)arytenoids and thyroid cartilages 168. Pneumoconiosis is caused by the 1)bacterium Streptococcus pneumoniae 2)inhalation of coal dust and sulphur particles 3)release of vasodilator substances due to allergens 4)deposition of iron particles in alveoli of lungs 169. The specialised myocytes present in inner ventricular walls of heart ben endocardium are called 1)Chorde tendinae 2)Papillary muscles 3)Purkinje fibres 4)Columnae carnea 170. In human heart, the non functional valve in the right atrium is present at the ope of 1)coronary sinus 2)superior venacava 3)inferior venacava 	167.	Rima glottidis is the	e narrow space betw	een the		
 168. Pneumoconiosis is caused by the bacterium Streptococcus pneumoniae inhalation of coal dust and sulphur particles release of vasodilator substances due to allergens deposition of iron particles in alveoli of lungs 169. The specialised myocytes present in inner ventricular walls of heart ben endocardium are called Chorde tendinae Papillary muscles Purkinje fibres Columnae carnea 170. In human heart, the non functional valve in the right atrium is present at the ope of coronary sinus superior venacava right atrioventricular aperture 		1)true vocal folds and	d false vocal cords	2)true vocal co	ords and arytenoids	
 1)bacterium <i>Streptococcus pneumoniae</i> 2)inhalation of coal dust and sulphur particles 3)release of vasodilator substances due to allergens 4)deposition of iron particles in alveoli of lungs 169. The specialised myocytes present in inner ventricular walls of heart ben endocardium are called 1)Chorde tendinae 2)Papillary muscles 3)Purkinje fibres 4)Columnae carnea 170. In human heart, the non functional valve in the right atrium is present at the ope of 1)coronary sinus 2)superior venacava 3)inferior venacava 		3)corniculate cartilag	ges and cuneiform cart	ilages 4)arytenoids	and thyroid cartilages	
 2)inhalation of coal dust and sulphur particles 3)release of vasodilator substances due to allergens 4)deposition of iron particles in alveoli of lungs 169. The specialised myocytes present in inner ventricular walls of heart ben endocardium are called 1)Chorde tendinae 2)Papillary muscles 3)Purkinje fibres 4)Columnae carnea 170. In human heart, the non functional valve in the right atrium is present at the ope of 1)coronary sinus 2)superior venacava 3)inferior venacava 4)right atrioventricular aperture 	168.	Pneumoconiosis is o	caused by the			
 3)release of vasodilator substances due to allergens 4)deposition of iron particles in alveoli of lungs 169. The specialised myocytes present in inner ventricular walls of heart ben endocardium are called 1)Chorde tendinae 2)Papillary muscles 3)Purkinje fibres 4)Columnae carnea 170. In human heart, the non functional valve in the right atrium is present at the ope of 1)coronary sinus 2)superior venacava 3)inferior venacava 	1)bacterium Streptococcus pneumoniae					
 4)deposition of iron particles in alveoli of lungs 169. The specialised myocytes present in inner ventricular walls of heart ben endocardium are called 1)Chorde tendinae 2)Papillary muscles 3)Purkinje fibres 4)Columnae carnea 170. In human heart, the non functional valve in the right atrium is present at the ope of 1)coronary sinus 2)superior venacava 3)inferior venacava 	2)inhalation of coal dust and sulphur particles					
 169. The specialised myocytes present in inner ventricular walls of heart ben endocardium are called 1)Chorde tendinae 2)Papillary muscles 3)Purkinje fibres 4)Columnae carnea 170. In human heart, the non functional valve in the right atrium is present at the ope of 1)coronary sinus 2)superior venacava 3)inferior venacava 		3)release of vasodilator substances due to allergens				
endocardium are called 1)Chorde tendinae 2)Papillary muscles 3)Purkinje fibres 4)Columnae carnea 170. In human heart, the non functional valve in the right atrium is present at the ope of 1)coronary sinus 2)superior venacava 3)inferior venacava 4)right atrioventricular aperture		4) deposition of iron	particles in alveoli of l	ungs		
1)Chorde tendinae2)Papillary muscles3)Purkinje fibres4)Columnae carnea170. In human heart, the non functional valve in the right atrium is present at the ope of01)coronary sinus2)superior venacava3)inferior venacava4)right atrioventricular aperture	169.	The specialised m	yocytes present in	inner ventricular	walls of heart beneat	
 170. In human heart, the non functional valve in the right atrium is present at the oper of 1)coronary sinus 3)inferior venacava 4)right atrioventricular aperture 		endocardium are ca	alled			
of1)coronary sinus2)superior venacava3)inferior venacava4)right atrioventricular aperture		1)Chorde tendinae	2)Papillary muscles	3)Purkinje fibres	4)Columnae carneae	
1)coronary sinus2)superior venacava3)inferior venacava4)right atrioventricular aperture	170.	In human heart, the non functional valve in the right atrium is present at the openin				
3)inferior venacava 4)right atrioventricular aperture		of				
		1)coronary sinus		2)superior venacava	a	
ROUCH		3)inferior venacava		4)right atrioventric	ular aperture	
ROUCH						
KOUGH			ROUG	GH		

171. Which of the following hormones are not antagonistic to each other in their actions?				
1)Parathormone and calcitonin	2)Somatocrinin and somatostatin			
3)Cortisol and Catecholamines	4)Aldosterone and atrial natriuretic peptide			
172. Prostaglandins are derivatives of				
1)tryptophan 2)tyrosine	3)glycoproteins 4)eicosanoids			
173. The vestibular apparatus concerned wi				
1)saccule and utricle	2)ampullae and cristae			
3)maculae and vestibule	4)semicircular canals and otolith organ			
174. Inter ventricular foramina are connect	-			
1)atria and ventricles in heart	2)paracoels and diocoel in brain			
3)3rd and 4th ventricles of brain	4)right and left ventricles of heart			
175. Joint between two adjacent vertebrae o	•			
1)symphysis and synarthrosis	2)synchondrosis and syndesmosis			
3)diarthrose and syndesmosis	4)symphysis and amphiarthrosis			
176. Hensen's disc is	, j I Januar I an and			
1)part of primitive streak	2)part of sarcomere made of only myosin			
3) primitive knot through which chordame				
4)bilaminar embryonic disc with epiblast a				
177. Autoimmune disease from the followin				
1)Glomerulo nephritis	2)Osteo arthritis			
3)Rheumatoid arthritis	4)Meningitis			
178. In the conduction of nerve impulse, relative refractory period coincides with the phase				
of				
1)depolarisation 2)repolarisation	3)hyperpolarisation 4)polarisation			
179. In kidneys, renal columns are made of				
1)renal calyces	2)medullary pyramids			
3)renal cortex	4)renal papillae			
180. In the process of urine formation, selective reabsorption is minimum in this part of				
nephron				
1)Proximal convoluted tubule	2)Distal convoluted tubule			
3)Henle's loop	4)Collecting duct			
ROUGH				